

BROWNIAN MOTION IN PLANETARY MIGRATION

RUTH A. MURRAY-CLAY¹ & EUGENE I. CHIANG^{1,2}

ACCEPTED TO APJ: July 10, 2006

ABSTRACT

A residual planetesimal disk of mass $10\text{--}100M_{\oplus}$ remained in the outer solar system following the birth of the giant planets, as implied by the existence of the Oort cloud, coagulation requirements for Pluto, and inefficiencies in planet formation. Upon gravitationally scattering planetesimal debris, planets migrate. Orbital migration can lead to resonance capture, as evidenced here in the Kuiper and asteroid belts, and abroad in extra-solar systems. Finite sizes of planetesimals render migration stochastic (“noisy”). At fixed disk mass, larger (fewer) planetesimals generate more noise. Extreme noise defeats resonance capture. We employ order-of-magnitude physics to construct an analytic theory for how a planet’s orbital semi-major axis fluctuates in response to random planetesimal scatterings. The degree of stochasticity depends not only on the sizes of planetesimals, but also on their orbital elements. We identify the conditions under which the planet’s migration is maximally noisy. To retain a body in resonance, the planet’s semi-major axis must not random walk a distance greater than the resonant libration width. We translate this criterion into an analytic formula for the retention efficiency of the resonance as a function of system parameters, including planetesimal size. We verify our results with tailored numerical simulations. Application of our theory reveals that capture of Resonant Kuiper belt objects by a migrating Neptune remains effective if the bulk of the primordial disk was locked in bodies having sizes $< \mathcal{O}(100)$ km and if the fraction of disk mass in objects with sizes $\gtrsim 1000$ km was less than a few percent. Coagulation simulations produce a size distribution of primordial planetesimals that easily satisfies these constraints. We conclude that stochasticity did not interfere with nor modify in any substantive way Neptune’s ability to capture and retain Resonant Kuiper belt objects during its migration.

Subject headings: celestial mechanics—Kuiper belt—diffusion—planets and satellites: formation—solar system: formation

1. INTRODUCTION

Planet formation by coagulation of planetesimals is not perfectly efficient—it leaves behind a residual disk of solids. Upon their coalescence, the outer planets of our solar system were likely embedded in a $10\text{--}100M_{\oplus}$ disk of rock and ice containing the precursors of the Oort cloud (Dones et al. 2004) and the Kuiper belt (see the reviews by Chiang et al. 2006; Cruikshank et al. 2006; Levison et al. 2006). The gravitational back-reaction felt by planets as they scatter and scour planetesimals causes the planets to migrate (Fernández & Ip 1984; Murray et al. 1998; Hahn & Malhotra 1999; Gomes, Morbidelli, & Levison 2004). Neptune is thought to have migrated outward and thereby trapped Kuiper belt objects (KBOs) into its exterior mean-motion resonances, both of low-order such as the 3:2 (Malhotra 1995) and of high-order such as the 5:2 (Chiang et al. 2003; Hahn & Malhotra 2005). Likewise, Jupiter’s inward migration may explain the existence of Hilda asteroids in 2:3 resonance with the gas giant (Franklin et al. 2004). A few pairs of extra-solar planets, locked today in 2:1 resonance (Vogt et al. 2005; Lee et al. 2006), may have migrated to their current locations within parent disks composed of gas and/or planetesimals. Orbital migration and resonant trapping of dust grains may also be required to explain non-axisymmetric

structures observed in debris disks surrounding stars $10\text{--}100$ Myr old (e.g., Wyatt 2003; Meyer et al. 2006).

Only when orbital migration is sufficiently smooth and slow can resonances trap bodies. The slowness criterion requires migration to be adiabatic: Over the time the planet takes to migrate across the width of the resonance, its resonant partner must complete at least a few librations. Otherwise the bodies speed past resonance (e.g., Dermott, Malhotra, & Murray 1988; Chiang 2003; Quillen 2006). Smoothness requires that changes in the planet’s orbit which are incoherent over timescales shorter than the libration time do not accumulate unduly. Orbital migration driven by gravitational scattering of discrete planetesimals is intrinsically not smooth. A longstanding concern has been whether Neptune’s migration was too “noisy” to permit resonance capture and retention (see, e.g., Morbidelli, Brown, & Levison 2003). In N-body simulations of migration within planetesimal disks (Hahn & Malhotra 1999; Gomes et al. 2004; Tsiganis et al. 2005), $N \sim \mathcal{O}(10^4)$ is still too small to produce the large, order-unity capture efficiencies seemingly demanded by the current census of Resonant KBOs.

At the same time, the impediment against resonance capture introduced by inherent stochasticity has been exploited to explain certain puzzling features of the Kuiper belt, most notably the Classical (non-Resonant) belt’s outer truncation radius, assumed to lie at a heliocentric distance of ~ 48 AU (Trujillo & Brown 2001; Levison & Morbidelli 2003). If Neptune’s 2:1 resonance captured KBOs and released them en route, Classical KBOs could have been transported (“combed”) outwards to popu-

¹ Center for Integrative Planetary Sciences, Astronomy Department, University of California at Berkeley, Berkeley, CA 94720, USA

² Alfred P. Sloan Research Fellow
 Electronic address: rmurray@astron.berkeley.edu, echiang@astron

late the space interior to the final position of the 2:1 resonance, at a semi-major axis of 47.8 AU (Levison & Morbidelli 2003). As originally envisioned, this scenario requires that $\sim 3M_{\oplus}$ be trapped inside the 2:1 resonance so that an attendant secular resonance suppresses growth of eccentricity during transport. It further requires that the degree of stochasticity be such that the migration is neither too smooth nor too noisy. Whether these requirements were actually met remain open questions.³

Stochastic migration has also been studied in gas disks, in which noise is driven by density fluctuations in turbulent gas. Laughlin, Steinacker, & Adams (2004) and Nelson (2005) propose that stochasticity arising from gas that is unstable to the magneto-rotational instability (MRI) can significantly prolong a planet’s survival time against accretion onto the parent star. The spectrum of density fluctuations is computed by numerical simulations of assumed turbulent gas.

In this work, we study stochastic changes to a planet’s orbit due to planetesimal scatterings. The planet’s Brownian motion arises from both Poisson variations in the rate at which a planet encounters planetesimals, and from random fluctuations in the mix of planet-planetesimal encounter geometries. How does the vigor of a planet’s random walk depend on the masses and orbital properties of surrounding planetesimals? We answer this question in §2 by constructing an analytic theory for how a migrating planet’s semi-major axis fluctuates about its mean value. We employ order-of-magnitude physics, verifying our assertions whenever feasible by tailored numerical integrations. Because the properties of planetesimal disks during the era of planetary migration are so uncertain, we consider a wide variety of possibilities for how planetesimal semi-major axes and eccentricities are distributed. One of the fruits of our labors will be identification of the conditions under which a planet’s migration is maximally stochastic.

Apportioning a fixed disk mass to fewer, larger planetesimals renders migration more noisy. How noisy is too noisy for resonance capture? What limits can we place on the sizes of planetesimals that would keep capture of Resonant KBOs by a migrating Neptune a viable hypothesis? These questions are answered in §3, where we write down a simple analytic formula for the retention efficiency of a resonance as a function of disk properties, including planetesimal size. Quantifying the size spectrum of planetesimals is crucial for deciphering the history of planetary systems. Many scenarios for the evolution of the Kuiper belt implicitly assume that most of the mass of the primordial outer solar system was locked in planetesimals having sizes of $\mathcal{O}(100)$ km, like those observed today (see, e.g., Chiang et al. 2006 for a critique of these scenarios). By contrast, coagulation simulations place the bulk of the mass in bodies having sizes of $\mathcal{O}(1)$ km (Kenyon & Luu 1999). For ice giant formation to proceed *in situ* in a timely manner in the outer solar sys-

tem, most of the primordial disk may have to reside in small, sub-km bodies (Goldreich, Lithwick, & Sari 2004).

In §4, in addition to summarizing our findings, we extend them in a few directions. The main thrust of this paper is to analyze how numerous, small perturbations to a planet’s orbit accumulate. We extend our analysis in §4 to quantify the circumstances under which a single kick to the planet from an extremely large planetesimal can disrupt the resonance. We also examine perturbations exerted directly on Resonant KBOs by ambient planetesimals.

2. STOCHASTIC MIGRATION: AN ORDER-OF-MAGNITUDE THEORY

We assume the planet’s eccentricity is negligibly small. We decompose the rate of change of the planet’s semi-major axis, $\dot{a}_p$, into average and random components,

$$\dot{a}_p = \dot{a}_{p,\text{avg}} + \dot{a}_{p,\text{rnd}}. \quad (1)$$

The average component (“signal”) arises from any global asymmetry in the way a planet scatters planetesimals, e.g., an asymmetry due to systematic differences between planetesimals inside and outside a planet’s orbit. The random component (“noise”) results from chance variations in the numbers and orbital elements of planetesimals interacting with the planet. By definition, $\dot{a}_{p,\text{rnd}}$ time-averages to zero. We assume that $\dot{a}_{p,\text{avg}}(t)$ is a known function of time t , and devote all of §2 to the derivation of $\dot{a}_{p,\text{rnd}}$.

Each close encounter between the planet and a single planetesimal lasting time Δt_e causes the planet’s semi-major axis to change by Δa_p . Expressions for Δa_p and Δt_e depend on the planetesimal’s orbital elements. We define $x \equiv a - a_p$ as the difference between the semi-major axes of the planetesimal and of the planet, $b > 0$ as the impact parameter of the encounter, and $u \sim e\Omega a$ as the planetesimal’s random (epicyclic) velocity, where a , e , and Ω are the semi-major axis, eccentricity, and mean angular velocity of the planetesimal, respectively. We assume that $|x| \lesssim a_p$. Encounters unfold differently according to how $|x|$ and b compare with the planet’s Hill radius,

$$R_H = a_p \left(\frac{M_p}{3M_*} \right)^{1/3}, \quad (2)$$

and according to how u compares with the Hill velocity,

$$v_H \equiv \Omega_p R_H. \quad (3)$$

Here M_p and M_* are the masses of the planet and of the star, respectively, and Ω_p is the angular velocity of the planet. See Table A1 for a listing of frequently used symbols.

In §2.1, we calculate Δa_p for a single encounter with a planetesimal having $|x| \gtrsim R_H$. In §2.2, we repeat the calculation for $|x| \lesssim R_H$. In §2.3, we provide formulae for the root-mean-squared (RMS) random velocity due to cumulative encounters, $\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2}$, and identify which cases of those treated in §§2.1–2.2 potentially yield the strongest degree of stochasticity in the planet’s migration.

³ While Classical KBOs do have semi-major axes that extend up to 48 AU, the distribution of their perihelion distances cuts off sharply at distances closer to 45 AU (see, e.g., Figure 2 of Chiang et al. 2006). Interpreted naively (i.e., without statistics), the absence of bodies having perihelion distances of 45–48 AU and eccentricities less than ~ 0.1 smacks of observational bias and motivates us to revisit the problem of whether an edge actually exists, or at least whether the edge bears any relation to the 2:1 resonance.

2.1. Single Encounters with $|x| \gtrsim R_H$: Non-Horseshoes

We calculate the change in the planet's semi-major axis, Δa_p , resulting from an encounter with a single planetesimal having $|x| \gtrsim R_H$. We treat planetesimals on orbits that do not cross that of the planet in §2.1.1 and those that do cross in §2.1.2. Throughout, Δ refers to the change in a quantity over a single encounter, evaluated between times well before and well after the encounter.

2.1.1. Non-Crossing Orbits

Planetesimals on orbits that do not cross that of the planet have

$$|x| > ae, \quad (4)$$

which corresponds to

$$|x|/R_H > u/v_H. \quad (5)$$

Our plan is to relate Δa_p to Δx by conservation of energy, calculate Δe using the impulse approximation, and finally generate Δx from Δe by conservation of the Jacobi integral.

By conservation of energy,

$$\Delta \left[-\frac{GM_* M_p}{2a_p} - \frac{GM_* m}{2a} \right] = 0, \quad (6)$$

where $m \ll M_p$ is the mass of the planetesimal. We have dropped terms that account for the potential energies of the planet and of the planetesimal in the gravitational field of the ambient disk. These are small because the disk mass is of order $M_p \ll M_*$ and because the disk does not act as a point mass but is spatially distributed. Equation (6) implies

$$\Delta a_p \sim -\frac{m}{M_p} \left(\frac{a_p}{a} \right)^2 \Delta a. \quad (7)$$

Since $|\Delta a_p| \ll |\Delta a|$ and $a_p \sim a$, we have $\Delta x \sim \Delta a$ and

$$\Delta a_p \sim -\frac{m}{M_p} \Delta x. \quad (8)$$

The impulse imparted by the planet changes the eccentricity of the planetesimal by Δe . An encounter for which $|x|$ is more than a few times R_H imparts an impulse per mass⁴

$$\Delta u \sim \pm \frac{GM_p}{b^2} \Delta t_e. \quad (9)$$

The impact parameter b is limited by

$$|x| - ae \lesssim b \lesssim |x| + ae. \quad (10)$$

Since $ae < |x|$,

$$b \sim |x|. \quad (11)$$

Because the relative speed due to Keplerian shear, $(3/2)\Omega_p|x|$, is larger than u , the relative speed during encounter is dominated by the former, and

$$\Delta t_e \sim \frac{2b}{(3/2)\Omega_p b} = \frac{4}{3\Omega_p} \sim \frac{1}{\Omega_p}. \quad (12)$$

Since Δt_e is about one-fifth of an orbital period, the impulse approximation embodied in (9) should yield good

⁴ The impulse to the planetesimal changes both u and the planetesimal's Keplerian shearing velocity, $-(3/2)\Omega x$. In the non-crossing case, $|\Delta u| > |\Delta(\Omega x)|$.

order-of-magnitude results. The change in the eccentricity of the planetesimal is hence

$$\Delta e \sim \frac{\Delta u}{\Omega_p a_p} \sim \pm \frac{M_p}{M_*} \left(\frac{a_p}{x} \right)^2. \quad (13)$$

When $|\Delta e| < e$ (the pre-encounter e), the change Δe can be either positive or negative, depending on the true anomaly of the planetesimal at the time of encounter. If $|\Delta e| > e$, then $\Delta e > 0$. When $|x| \sim R_H$, $|\Delta e|$ attains its maximum value of $\sim (M_p/M_*)^{1/3}$; i.e., $\Delta u \sim v_H$.⁵

To calculate the corresponding change in the planetesimal's semi-major axis, Δx , we exploit conservation of the Jacobi integral, C_J . That a conserved integral exists relies on the assumption that in the frame rotating with the planet, the potential (having a centrifugal term plus gravitational contributions due to the star, planet, and disk) is time-stationary; the Jacobi integral is simply the energy of the planetesimal (test particle) evaluated in that frame. To the same approximation embodied in Equation (6),

$$\begin{aligned} -\frac{1}{2}C_J &= E - \Omega_p J \\ &= -\frac{GM_*}{a_p} \left[\frac{1}{2(a/a_p)} + \sqrt{(a/a_p)(1-e^2)} \right] \end{aligned} \quad (14)$$

far from encounter, where E and J are the energy and angular momentum per mass of the planetesimal, respectively. Taking the differential of (14) yields, to leading order,

$$\frac{3}{4} \left(\frac{\Delta x}{a_p} \right)^2 + \left(\frac{3}{2} \frac{x}{a_p} - \frac{1}{2} e^2 \right) \frac{\Delta x}{a_p} - \Delta(e^2) = 0. \quad (15)$$

Since $|x|/a_p > e > e^2$ (non-crossing condition) and $\Delta(e^2) < (x/a_p)^2$ (by Equation [13] and the condition $|x| > R_H$), Equation (15) reduces to

$$\Delta x \sim \frac{2a_p^2}{3x} \Delta(e^2). \quad (16)$$

We combine Equations (8) and (16) to find

$$\Delta a_p \sim -\frac{m}{M_p} \frac{a_p^2}{x} \Delta(e^2). \quad (17)$$

Equation (17) takes two forms depending on how $|\Delta e|$ compares with (the pre-encounter) e . If

$$|x| > R_H \left(\frac{v_H}{u} \right)^{1/2}, \quad (18)$$

then $|\Delta e| < e$, $\Delta(e^2) \sim 2e\Delta e$, and Equation (17) becomes

$$\Delta a_p \sim \mp \frac{m}{M_*} \frac{a_p^4}{x^3} e. \quad (19)$$

⁵ When $|x| \lesssim 2R_H$, the encounter pulls the planetesimal into the planet's Hill sphere. The planetesimal accelerates in a complicated way and exits the Hill sphere in a random direction with u of order the planet's escape velocity at the Hill radius, v_H (Petit & Hénon 1986). The planetesimal's eccentricity is boosted by $\Delta e \sim (M_p/M_*)^{1/3}$. The encounter time is typically the time required to complete a few orbits around the planet, $\Delta t_e \sim 2\pi/\Omega_p$. Since Δt_e and Δe match, to order of magnitude, Equations (12) and (13) for $|x| \sim R_H$, we do not treat $R_H \lesssim |x| \lesssim 2R_H$ as an explicitly different case.

The right-hand side is extremized for $|x| \sim R_H (v_H/u)^{1/2}$:

$$\max |\Delta a_p| \sim \frac{m}{M_p} R_H \left(\frac{a_p e}{R_H} \right)^{5/2} < \frac{m}{M_p} R_H, \quad (20)$$

valid for $e < R_H/a_p$ (non-crossing).

On the other hand, if

$$R_H \lesssim x < R_H \left(\frac{v_H}{u} \right)^{1/2}, \quad (21)$$

then $|\Delta e| > e$ and $\Delta(e^2) \sim (\Delta e)^2$, and Equation (17) becomes

$$\Delta a_p \sim -\frac{m M_p}{M_*^2} \frac{a_p^6}{x^5}. \quad (22)$$

Equation (22) agrees with the more careful solution of Hill's problem by Hénon & Petit (1986). The right-hand side is extremized for $|x| \sim R_H$:

$$\max |\Delta a_p| \sim \frac{m}{M_p} R_H. \quad (23)$$

In summary, if $x/R_H > u/v_H$, then (a) the planetesimal's orbit does not cross ("nc") that of the planet, (b)

$$\Delta t_e \sim \frac{1}{\Omega_p}, \quad (24)$$

and (c)

$$\Delta a_p = \begin{cases} \Delta a_{p,nc1} \sim -\frac{m M_p}{M_*^2} \frac{a_p^6}{x^5}, & \text{if } R_H \lesssim x \lesssim (v_H/u)^{1/2} R_H; \\ \Delta a_{p,nc2} \sim \mp \frac{m}{M_*} \frac{a_p^4}{x^3} e, & \text{if } x \gtrsim (v_H/u)^{1/2} R_H. \end{cases} \quad (25)$$

2.1.2. Crossing Orbits

Encounters with planetesimals on orbits that cross that of the planet, i.e., those with

$$|x|/R_H < u/v_H, \quad (26)$$

differ from encounters with non-crossing planetesimals in two key respects. First, the relative velocity of the two bodies is dominated by the planetesimal's random (epicyclic) velocity rather than the Keplerian shear. Second, the planetesimal's impact parameter, b , may differ significantly from $|x|$. The impact parameter may take any value

$$b_{\min} < b \lesssim a e, \quad (27)$$

where $b_{\min}$ is the impact parameter below which the planetesimal collides with the planet. Because crossing orbits allow for encounters with many different geometries, outcomes of these encounters can vary dramatically. Here we restrict ourselves to estimating the maximum $|\Delta a_p|$ that can result from an orbit-crossing encounter. In §2.3.2, we argue this restriction is sufficient for our purposes.

When $u > v_H$, the eccentricity of the planetesimal can change by at most $|\Delta e| \sim e$. Such a change corresponds to an order-unity rotation of the direction of the planetesimal's random velocity vector, and requires that

$b \lesssim G M_p / u^2$. The change in the planetesimal's specific energy over the encounter is approximately

$$\Delta \left(-\frac{G M_*}{2a} \right) \sim \Delta \left(\frac{1}{2} v^2 \right) + \Delta \left(-\frac{G M_*}{r} \right), \quad (28)$$

where v is the velocity of the planetesimal relative to the star (in an inertial frame of reference) and r is the distance between the planetesimal and the star. Now Δr for an encounter with $b \lesssim G M_p / u^2$ is at most $G M_p / u^2 < R_H$ and $\Delta(v^2)$ is of order $\Omega a u$. Since

$$\Omega a u \gtrsim \Omega^2 a R_H > \Omega^2 a \frac{G M_p}{u^2}, \quad (29)$$

the second term on the right-hand side of Equation (28) is negligible compared to the first, and the maximum $|\Delta a|$ over an encounter is

$$\max |\Delta a| \sim \frac{a^2}{G M_*} \Omega a u \sim a e. \quad (30)$$

By Equation (7) and $a \sim a_p$,

$$\max |\Delta a_p| \sim \frac{m}{M_p} a_p e. \quad (31)$$

We have verified Equation (31) by numerical orbit integrations. We could also have arrived at Equation (31) through Equation (15), which yields $|\Delta x| \sim a_p e$ for crossing orbits when $|\Delta e| \sim e$.

2.2. Single Encounters with $|x| \lesssim R_H$: Horseshoes

When $|x| < R_H$, planetesimals can occupy horseshoe orbits. A planetesimal on a horseshoe orbit for which $|x| \approx R_H$ encounters the planet on a timescale somewhat shorter than the orbital period; by the impulse approximation, such a planetesimal kicks the planet such that

$$\Delta a_p \sim \frac{m}{M_*} \frac{a_p^3}{b^2}, \quad (32)$$

where we have momentarily restricted consideration to planetesimals having sub-Hill eccentricities ($e \lesssim R_H/a_p$). From Hénon and Petit (1986),

$$b = \frac{8}{3} \frac{R_H^3}{x^2}, \quad (33)$$

valid for x not too far below R_H . Then

$$|\Delta a_p| \sim \frac{m}{M_p} \frac{x^4}{R_H^3}. \quad (34)$$

The kick is maximal for maximum $|x| = R_H$:

$$\max |\Delta a_p| \sim \frac{m}{M_p} R_H. \quad (35)$$

This is the same maximum as was derived for the $|x| \sim R_H$, non-crossing case; see Equation (23). Thus, a co-orbital ring of planetesimals on horseshoe orbits with sub-Hill eccentricities increases the stochasticity generated by planetesimals on non-horseshoe, non-crossing orbits by a factor of at most order unity (under the assumption that disk properties are roughly constant within several Hill radii of the planet). For this reason, and also because the horseshoe region may well have been depleted of planetesimals compared to the rest of the disk, we omit consideration of co-orbital, sub-Hill planetesimals

for the remainder of the paper, confident that the error so incurred will be at most order unity.

What about planetesimals on horseshoe orbits with super-Hill eccentricities ($e \gtrsim R_H/a_p$)? Upon encountering the planet, such objects can have their semi-major axes changed by $|\Delta a| > R_H$ —whereupon they are expelled from the 1:1 horseshoe resonance. Because highly eccentric, horseshoe resonators are unstable, we neglect consideration of them for the rest of our study.

2.3. Multiple Encounters: Cumulative Stochasticity

We now extend our analysis from individual encounters to the cumulative stochasticity generated by a disk with surface density Σ_m in planetesimals of a single mass m . Note that Σ_m need not equal the total surface density Σ (integrated over all possible masses m). We will consider size distributions in §3.6. We consider planetesimals with sub-Hill ($u < v_H$) velocities (§2.3.1) separately from those with super-Hill ($u > v_H$) velocities (§2.3.2). Sub-Hill (non-horseshoe) planetesimals always occupy non-crossing orbits. Super-Hill planetesimals can be crossing or non-crossing.

2.3.1. Sub-Hill Velocities ($u < v_H$)

Consider planetesimals with sub-Hill velocities located a radial distance x away from the planet ($|x| > R_H$). Since $u < v_H$, the speeds of planetesimals relative to the planet are determined principally by Keplerian shear (Equation [12]), and the scale height of the planetesimals is less than R_H . The planet encounters (undergoes conjunctions with) such planetesimals at a mean rate

$$\dot{N} \sim \frac{\Sigma_m}{m} \Omega x^2, \quad (36)$$

as is appropriate for encounters in a two-dimensional geometry. Over a time interval Δt , the planet encounters $\bar{N} = \dot{N} \Delta t$ such planetesimals on average. Systematic trends in $\bar{N}$ with x —say, systematically more objects encountered interior to the planet’s orbit than exterior to it—cause the planet to migrate along an average trajectory with velocity $\dot{a}_{p,\text{avg}}$.

Random fluctuations in (a) the number of planetesimals encountered per fixed time interval and (b) the mix of planetesimals’ pre-encounter orbital elements cause the planet to random walk about this average trajectory. Contribution (a) is straightforward to model. The probability that the planet encounters N objects located a distance x away in time Δt is given by Poisson statistics:

$$P(N) = \frac{\bar{N}^N}{N!} e^{-\bar{N}}. \quad (37)$$

The variance in N is

$$\sigma_N^2 \equiv \langle (N - \bar{N})^2 \rangle = \bar{N}. \quad (38)$$

Fluctuations in N drive the planet either towards or away from the star with equal probability and with typical speed

$$\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2} \sim \frac{|\Delta a_p|}{\Delta t} \bar{N}^{1/2}, \quad (39)$$

hereafter the root-mean-squared (RMS) speed. While $\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2} \propto 1/\sqrt{\Delta t}$, the distance random walked $\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2} \Delta t \propto \sqrt{\Delta t}$.

Our assumption of Poisson statistics is reasonable. In the sub-Hill case, a planet-planetesimal encounter requires a time $\Delta t_e \sim 1/\Omega_p$ (Equation [12]) to complete. Encounters separated by more than Δt_e are uncorrelated with one another, at least until the planet completes one revolution with respect to the surrounding disk, i.e., at least until a synodic time $t_{\text{syn}} \sim 4\pi a_p / (3\Omega_p |x|)$ elapses. After a synodic period, it is possible, in principle, for the planet to essentially repeat the same sequence of encounters that it underwent during the last synodic period. We assume in this paper that this does not happen—that the orbits of planetesimals interacting with the planet are randomized on a timescale $t_{\text{rdz}} < t_{\text{syn}}$. We expect this inequality to be enforced by a combination of (i) randomization of planetesimal orbits due to encounters with the planet (e.g., encounters within the chaotic zone of the planet [Wisdom 1980]), (ii) phase mixing of planetesimals due to Keplerian shear (which occurs on timescale t_{syn} for planetesimals distributed between x and $\sim 2x$), (iii) gravitational interactions between planetesimals, and (iv) physical collisions between planetesimals. As long as $t_{\text{rdz}} < t_{\text{syn}}$, we are free to choose Δt to be anything longer than Δt_e .⁶

Contribution (b) is difficult to model precisely since we do not know how orbital elements of planetesimals are distributed. These distributions are unlikely to be governed by simple Poisson or Gaussian statistics (see, e.g., Ida & Makino 1992; Rafikov 2003; Collins & Sari 2006). Nonetheless, neglecting contribution (b) will not lead to serious error. Suppose the planetesimals’ pre-encounter elements are distributed such that the fractional variation in each element is at most of order unity (e.g., the planetesimal eccentricities span a range from $e/2$ to $2e$ at most). Then the central limit theorem ensures that the noise introduced by random sampling of orbital elements is at most comparable to the noise introduced by random fluctuations in the encounter rate. Consider, for example, noise that arises from random sampling of e in the case where $\Delta a_p = \Delta a_{p,\text{nc2}}$ (Equation [25]). For an encounter rate fixed at $\dot{N}$, the planet’s semi-major axis a_p changes over time interval Δt by $\bar{N} \times \overline{\Delta a_p}$, where $\overline{\Delta a_p}$ is the mean of $\bar{N} = \dot{N} \Delta t$ sampled values of Δa_p . If the dispersion in e for individual planetesimals is σ_e and the mean eccentricity sampled over $\bar{N}$ values is $\bar{e}$, then the dispersion in the sampled mean eccentricity is $\sigma_{\bar{e}} \sim \sigma_e / \bar{N}^{1/2}$ by the central limit theorem. For $\Delta a_p = \Delta a_{p,\text{nc2}} \propto e$, the dispersion in $\overline{\Delta a_p}$ is $|\Delta a_p| \sigma_{\bar{e}} / \bar{e}$. The planet’s RMS speed generated purely from random sampling of e is

$$\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2} \sim \frac{\bar{N}}{\Delta t} |\Delta a_p| \frac{\sigma_{\bar{e}}}{\bar{e}} \sim \frac{|\Delta a_p|}{\Delta t} \bar{N}^{1/2} \frac{\sigma_e}{\bar{e}}, \quad (40)$$

which is at most comparable to the RMS speed generated purely from random sampling of N (Equation [39]),

⁶ If $t_{\text{rdz}} > t_{\text{syn}}$, then $\Delta t > t_{\text{rdz}}$ and the right-hand side of Equation (39) is multiplied by $\sqrt{t_{\text{rdz}}/t_{\text{syn}}}$. The planet’s motion is more stochastic in this case because over t_{rdz} , correlated interactions with planetesimals do not cancel each other as much as uncorrelated interactions would. Later, since we will be interested in stochastic perturbations to mean-motion resonant particles, we will require $t_{\text{rdz}} < t_{\text{lib}}$, where t_{lib} is the libration period within resonance.

as desired. Our original supposition that $\max(\sigma_e/e) \sim 1$ leads to no important loss of generality; if the distribution in e were bi-modal, for example, we could treat each population separately and add the resultant RMS speeds in quadrature. Similar results obtain for random sampling of other elements such as x . For simplicity, we hereafter treat explicitly only fluctuations in the encounter rate [Equation (39)], knowing that the noise so calculated will be underestimated by a factor of at most order unity.

Expression (39) measures the contribution to the RMS speed from planetesimals located a distance x away. Since Δa_p scales inversely with x to a steep power in the sub-Hill regime (see [25]), the contribution to the RMS speed is greatest from objects at small x (for reasonable variations of Σ_m with x). We take disk material to extend to a minimum distance of $|x_{\min}| \equiv \mathcal{R}R_H$ ($\mathcal{R} > 1$) from the planet's orbit. Insertion of (25) into (39) yields

$$\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2} \sim \begin{cases} \mathcal{R}^{-4} \frac{1}{(\Omega_p \Delta t)^{1/2}} \left(\frac{\Sigma_m a_p^2 m}{M_p^2} \right)^{1/2} \frac{R_H}{a_p} v_H, & \text{if } 1 < \mathcal{R} < (v_H/u)^{1/2}; \\ \mathcal{R}^{-2} \frac{1}{(\Omega_p \Delta t)^{1/2}} \left(\frac{\Sigma_m a_p^2 m}{M_p^2} \right)^{1/2} e v_H, & \text{if } \mathcal{R} > (v_H/u)^{1/2}. \end{cases} \quad (41)$$

The RMS speed is maximized for $\mathcal{R} = 1$.

2.3.2. Super-Hill Velocities ($u > v_H$)

Next we consider the noise generated by planetesimals with super-Hill random velocities ($u > v_H$). We refer to close encounters that change a_p by the maximal amount, $\max|\Delta a_p| \sim (m/M_p)a_p e$ (Equation [31]), as “maximal encounters.” Maximal encounters, which occur at impact parameters $b \lesssim GM_p/u^2$, make an order-unity contribution to the total super-Hill stochasticity. Non-maximal (more distant) encounters contribute to the total stochasticity through a Coulomb-like logarithm, as we show at the end of this sub-section.

For a maximal encounter, a planetesimal must approach within distance $b \lesssim GM_p/u^2$. Such encounters occur at a mean rate

$$\dot{N} \sim n \left(\frac{GM_p}{u^2} \right)^2 u \sim \frac{\Sigma_m}{m} \Omega_p R_H^2 \left(\frac{v_H}{u} \right)^4, \quad (42)$$

where $n \sim \Sigma_m \Omega / (mu)$ is the number density of planetesimals, and we have assumed that planetesimal inclinations and eccentricities are of the same order. Over a time interval Δt , the planet encounters on average $\dot{N} \Delta t$ such planetesimals, each of which increases or decreases a_p by about $\max|\Delta a_p|$. Since planetesimals suffering maximal encounters have their orbits effectively randomized relative to each other, we may choose Δt to be any time interval longer than an encounter time $\Delta t_e \sim b/u < 1/\Omega_p$ (see related discussion in §2.3.1). Therefore the planet random walks with RMS velocity (averaged over time Δt)

$$\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2} \sim \frac{\dot{N} \Delta t^{1/2} \max|\Delta a_p|}{\Delta t}$$

$$\sim \left(\frac{1}{\Omega_p \Delta t} \right)^{1/2} \left(\frac{\Sigma_m a_p^2 m}{M_p^2} \right)^{1/2} \frac{v_H}{u} \frac{R_H}{a_p} v_H. \quad (43)$$

What about the contribution from non-maximal encounters? For a super-Hill encounter at impact parameter b , the specific impulse imparted to the planetesimal is $\sim GM_p/(bu)$. We suppose that $|\Delta a_p|$ is proportional to this specific impulse, so that $|\Delta a_p| \propto 1/b$. We have confirmed this last proportionality by numerical orbit integrations (not shown). Since $\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2} \propto (\dot{N})^{1/2} |\Delta a_p|$ and $\dot{N} \propto b^2$, we have $\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2} \propto b^0$, which implies that each octave in impact parameter contributes equally to the total stochasticity. In other words, our estimate for $\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2}$ in Equation (43) should be enhanced by a logarithmic factor of $\ln(b_{\max}/b_{\min})$, where $b_{\max}$ and $b_{\min} \sim GM_p/u^2$ are maximum and minimum impact parameters. We estimate $b_{\max} \sim u/\Omega$, the value for which the relative velocity of a super-Hill encounter is dominated by the planetesimal's random velocity rather than by the background shear. The logarithm is not large; for example, for $e = 0.2$, $\ln[(u/\Omega)/(GM_p/u^2)] \sim 5$.

2.3.3. Summary

We can neatly summarize Equations (41) and (43) by defining the Hill eccentricity,

$$e_H \equiv R_H/a_p, \quad (44)$$

and parameterizing Σ_m such that the disk contains mass $\mathcal{M}M_p$ in planetesimals of mass m spread uniformly from $a_d/2$ to $3a_d/2$:

$$\Sigma_m = \frac{\mathcal{M}M_p}{2\pi a_d^2}, \quad (45)$$

where $\mathcal{M}$ is a dimensionless number of order unity. Then

$$\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2} \sim \begin{cases} \mathcal{C} \mathcal{R}^{-4} (\Omega_p \Delta t)^{-1/2} \left(\frac{\mathcal{M}m}{M_p} \right)^{1/2} \frac{a_p}{a_d} e_H v_H, & \text{if } e < e_H/\mathcal{R}^2; \\ \mathcal{C} \mathcal{R}^{-2} (\Omega_p \Delta t)^{-1/2} \left(\frac{\mathcal{M}m}{M_p} \right)^{1/2} \frac{a_p}{a_d} e v_H, & \text{if } e_H/\mathcal{R}^2 < e < \mathcal{R} e_H; \\ \mathcal{C} (\Omega_p \Delta t)^{-1/2} \left(\frac{\mathcal{M}m}{M_p} \right)^{1/2} \frac{a_p}{a_d} \frac{e_H^2}{e} v_H, & \text{if } e > \mathcal{R} e_H, \end{cases} \quad (46)$$

where we have introduced a constant coefficient $\mathcal{C}$ (the same for each case so that the function remains continuous across case boundaries). The coefficient $\mathcal{C}$ encapsulates all the factors of order unity that we have dropped in our derivations. By studying N-body simulation data pertaining to the case $e > \mathcal{R} e_H$ as recorded in the literature (Hahn & Malhotra 1999; Gomes et al. 2004), we estimate that $\mathcal{C}$ is possibly of the order of several. That $\mathcal{C} > 1$ (but not $\gg 1$) is consonant with our having consistently underestimated the noise by neglecting (a) distant, non-maximal encounters in the super-Hill

regime (§2.3.2), and (b) stochasticity introduced by random sampling of orbital elements of planetesimals encountering the planet (§2.3.1). We defer definitive calibration of $\mathcal{C}$ to future study, but retain the coefficient in our expressions below to assess the degree to which our quantitative estimates are uncertain.

Planetesimals with $e = \mathcal{R}e_H$ ($|x_{\min}| = a_p e$) produce the maximum possible stochasticity:

$$\max \langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2} \sim \frac{\mathcal{C}}{\mathcal{R}} (\Omega_p \Delta t)^{-1/2} \left(\frac{\mathcal{M}m}{M_p} \right)^{1/2} \frac{a_p}{a_d} e_H v_H, \quad \text{for } e = \mathcal{R}e_H \geq e_H. \quad (47)$$

We will often adopt this case for illustration purposes below.

Since $\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2} \propto (\mathcal{M}m)^{1/2}$, the stochasticity driven by planetesimals having a range of sizes is dominated by those objects (in some logarithmic size bin) having maximal $\Sigma_m m$. For common power-law size distributions, such objects will occupy the upper end of the distribution. We will explicitly consider various possible size distributions in §3.6.

3. APPLICATION: MAXIMUM PLANETESIMAL SIZES

Neptune is thought to have migrated outward by scattering planetesimals during the late stages of planet formation (Fernández & Ip 1984; Hahn & Malhotra 1999). As the planet migrated, it may have captured Kuiper belt objects into its exterior resonances (Malhotra 1995), giving rise to the Resonant KBOs observed today (Chiang et al. 2003; Hahn & Malhotra 2005). If Neptune's migration had been too stochastic, however, resonance capture could not have occurred. A planetesimal disk having fixed surface mass density Σ generates more stochasticity when composed of larger (fewer) planetesimals. Therefore, assuming that planetary migration and concomitant resonance capture correctly explain the origin of present-day Resonant KBOs, we can rule out size distributions that are too “top-heavy” during the era of migration.

Stochasticity causes a planet to migrate both outward and inward. In §3.1, we provide background information regarding how resonance capture and retention depend on the sign of migration. In §3.2, we lay out general considerations for whether a stochastically migrating planet can retain particles in resonance. In §3.3, we derive and evaluate analytic, order-of-magnitude expressions for the maximum planetesimal size compatible with resonance retention, in the simple case when all planetesimals have the same size. In §3.4, we provide an analytic formula that details precisely how the resonance retention efficiency varies with average migration speed and planetesimal size. These analytic results are quantitatively tested by numerical integrations in §3.5. Cases where planetesimals exhibit a wide range of sizes are examined in §3.6.

3.1. Migrating Outward and Inward

As a planet migrates smoothly outward (away from the parent star), it can capture planetesimals into its exterior mean-motion resonances. By contrast, a planet which migrates smoothly inward cannot capture planetesimals which are initially non-resonant into its exterior resonances (e.g., Peale 1986). But a planetesimal that starts in exterior resonance with an inwardly migrating planet can remain in resonance for a finite time.




FIG. 1.— Evolution of a planetesimal in external 3:2 resonance as the planet migrates smoothly inward. The planetesimal remains in resonance until its eccentricity reaches zero, at a semi-major axis of $a = a_{\min}$.

Goldreich (1965) demonstrates how an outwardly migrating body adds angular momentum to a test particle in exterior resonance at just the right rate to keep the particle in resonance. Reversing the signs in his proof implies that an inwardly migrating body removes angular momentum from a particle in exterior resonance, pulling it inward while preserving the resonant lock. A planetesimal's eccentricity decreases as it is pulled inward. The adiabatic invariant,

$$\mathcal{N} = \sqrt{GM_* a} (p\sqrt{1-e^2} - q), \quad (48)$$

which is preserved for migration timescales long compared to the synodic time (e.g., Murray-Clay & Chiang 2005), implies that a planetesimal in $p : q$ exterior resonance ($p > q$) cannot be pulled inward to a semi-major axis less than

$$a_{\min} = \frac{1}{GM_*} \left(\frac{\mathcal{N}}{p-q} \right)^2 = a_0 \left(\frac{p\sqrt{1-e_0^2} - q}{p-q} \right)^2, \quad (49)$$

the value for which $e = 0$. Here a_0 and e_0 are the initial semi-major axis and eccentricity of the planetesimal, respectively.

Thus an exterior particle follows an inwardly migrating planet in resonant lockstep until it either reaches zero eccentricity (view Figure 4 of Peale 1986 or Figure 8.22 of Murray & Dermott 1999 in reverse) or until it crosses the separatrix (view Figure 5 of Peale 1986 or Figure 8.23 of Murray & Dermott 1999 in reverse), whichever comes first. We illustrate the former possibility in Figure 1 and the latter possibility in Figure 2, using our own orbit integrations. The value of $a_{\min}$ is annotated for reference.




FIG. 2.— Evolution of a planetesimal in external 3:2 resonance as the planet migrates smoothly inward. The planetesimal remains in resonance until it crosses the separatrix; by contrast to the evolution shown in Figure 1, the particle’s semi-major axis does not reach $a_{\min}$ and its eccentricity stays greater than zero.

3.2. General Considerations for Resonance Retention

The random component of the planet’s migration is a form of Brownian motion. The planet encounters a large number of small planetesimals, each of which causes the planet’s semi-major axis a_p to randomly step a small distance. Each random change in a_p produces a corresponding change in the semi-major axis of exact resonance,⁷ but no corresponding change in the actual semi-major axis of a resonant particle. The semi-major axis of the particle does not respond because a_p changes randomly every encounter time Δt_e , which is much shorter than the resonant libration period. Only changes in a_p that are coherent over timescales longer than the libration period produce an adiabatic response in the particle’s semi-major axis. We have verified these assertions by numerical orbit integrations (not shown).

The no-response condition implies that as a_p random walks, the difference between the semi-major axes of exact resonance and of a resonant particle random walks correspondingly. In other words, a resonant particle’s libration amplitude random walks. The sign and magnitude of each step in the libration amplitude’s random walk depend on the phase of libration when the step is taken. Since at any given time an ensemble of resonant particles are distributed over the full range of phases, a single random-walk history for the planet generates an ensemble of different random-walk histories for the particles.

When the libration amplitude of a resonant particle

⁷ A particle in exact resonance has zero libration amplitude, by definition.

random walks past its maximum allowed value, the particle escapes resonance. The maximum libration amplitude (full width) as measured in semi-major axis is

$$\delta a_{p,\text{lib}} = 2C_{\text{lib}}a_p \left(\frac{M_p e_{\text{res}}}{M_*} \right)^{1/2}, \quad (50)$$

where e_{res} is the eccentricity of the resonant object (not to be confused with the planetesimals generating the bulk of the noise), $C_{\text{lib}} \approx 4\sqrt{f_{31}/3}$ is a constant (see Murray and Dermott 1999 for f_{31}), and we have restricted consideration to first-order ($p - q = 1$) resonances. For the 3:2 exterior resonance with Neptune, $C_{\text{lib}} \approx 3.64$. Note that in contrast to the usual definition of maximum libration width, $\delta a_{p,\text{lib}}$ refers not to the particle’s semi-major axis, but rather to the planet’s. The meaning of $\delta a_{p,\text{lib}}$ is as follows. Take a particle in exact resonance. By definition, such a particle has zero libration amplitude. Then the planet’s semi-major axis can change instantaneously by at most $\delta a_{p,\text{lib}}/2$ and the particle will still remain in resonance (but with finite libration amplitude).

Equation (50) derives from the pendulum model of resonance, which is known to be inaccurate at large e_{res} for some resonances. Malhotra (1996) finds numerically that for $e_{\text{res}} = 0.1$ – 0.4 , $\delta a_{p,\text{lib}}$ for the 3:2 resonance is insensitive to e_{res} , whereas the pendulum model predicts that $\delta a_{p,\text{lib}}$ doubles over this range. We nevertheless employ Equation (50) to estimate the maximum libration width, since it is simple, analytic, and introduces errors less than of order unity in our numerical evaluations below. The qualitative physics described in this paper does not depend on the accuracy to which we estimate $\delta a_{p,\text{lib}}$.

Consider a planet which migrates outward on average. When the random component of the planet’s migration is added to the average component, a planet can migrate either outward or inward at any moment. Call $S_{\text{rnd}} = \int_0^t \dot{a}_{p,\text{rnd}} dt$ the running sum of the random changes in a_p . The probability P_{keep} that a given particle is retained in resonance over some duration of migration equals the probability that $|S_{\text{rnd}}|$ remains less than the maximum libration half-width $\delta a_{p,\text{lib}}/2$ during that time. A particle that escapes resonance by being dropped behind the resonance ($S_{\text{rnd}} = +\delta a_{p,\text{lib}}/2$) is, practically speaking, permanently lost. The planet cannot recapture the particle by smoothly migrating inward (see §3.1). The random component of the planet’s migration can cause the planetesimal to be recaptured, but a recaptured particle lies on a trajectory near the separatrix and quickly re-escapes in practice. Once the average (outward) component of the planet’s migration carries the resonance well past the particle, the particle cannot be recaptured even if S_{rnd} random walks back to zero; in other words, the particle has been permanently left behind. A particle that escapes by being dropped in front of the resonance ($S_{\text{rnd}} = -\delta a_{p,\text{lib}}/2$) is also lost more often than not. Such a particle can be recaptured when the planet resumes migrating outward. Nevertheless, upon its recapture onto a trajectory near the separatrix, the particle can librate back to smaller semi-major axes and be expelled behind the resonance permanently.

3.3. Order-of-Magnitude Planetesimal Sizes

Armed with the considerations of §3.2, we are now ready to derive analytic, order-of-magnitude expressions

for the maximum planetesimal sizes compatible with resonance retention, for the simple case when the disk is composed of objects of a single size. The assumption of a single size is relaxed in §3.6.

Say the planet takes time T to migrate at speed $\dot{a}_{p,\text{avg}}$ from its initial to its final semi-major axis. Over this time, a_p random walks an expected distance of

$$\sigma_{a_p,T} \sim \langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2} T \sim \begin{cases} \mathcal{C} \mathcal{R}^{-4} \left(\frac{\mathcal{M}m}{M_p} \right)^{1/2} \frac{a_p}{a_d} e_H v_H \left(\frac{T}{\Omega_p} \right)^{1/2}, & \text{if } e < e_H/\mathcal{R}^2; \\ \mathcal{C} \mathcal{R}^{-2} \left(\frac{\mathcal{M}m}{M_p} \right)^{1/2} \frac{a_p}{a_d} e v_H \left(\frac{T}{\Omega_p} \right)^{1/2}, & \text{if } e_H/\mathcal{R}^2 < e < \mathcal{R}e_H; \\ \mathcal{C} \left(\frac{\mathcal{M}m}{M_p} \right)^{1/2} \frac{a_p}{a_d} \frac{e_H^2}{e} v_H \left(\frac{T}{\Omega_p} \right)^{1/2}, & \text{if } e > \mathcal{R}e_H, \end{cases} \quad (51)$$

where we have set $\Delta t = T$ in evaluating $\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2}$.

If $\sigma_{a_p,T} < \delta a_{p,\text{lib}}/2$, the planet can keep a large fraction of planetesimals in resonance. That is, most particles are retained in resonance when the disk mass comprises planetesimals of mass

$$m \lesssim m_{\text{crit}} \sim \begin{cases} \frac{\mathcal{R}^8 C_{\text{lib}}^2}{\mathcal{C}^2 \mathcal{M}} \left(\frac{a_d}{a_p} \right)^2 \frac{1}{\Omega_p T} \frac{e_{\text{res}}}{e_H} M_p, & \text{if } e < e_H/\mathcal{R}^2; \\ \frac{\mathcal{R}^4 C_{\text{lib}}^2}{\mathcal{C}^2 \mathcal{M}} \left(\frac{a_d}{a_p} \right)^2 \frac{1}{\Omega_p T} \frac{e_{\text{res}} e_H}{e^2} M_p, & \text{if } e_H/\mathcal{R}^2 < e < \mathcal{R}e_H; \\ \frac{C_{\text{lib}}^2}{\mathcal{C}^2 \mathcal{M}} \left(\frac{a_d}{a_p} \right)^2 \frac{1}{\Omega_p T} \frac{e_{\text{res}} e^2}{e_H^3} M_p, & \text{if } e > \mathcal{R}e_H. \end{cases} \quad (52)$$

Equation (52) can be equivalently interpreted as an upper limit on T for planetesimals of given mass m . For a fixed degree of noise, resonant objects are more difficult to retain if the average migration is slow.

We evaluate (52) to estimate the maximum planetesimal radius, $s = (3m/4\pi\rho)^{1/3}$, compatible with resonant capture of KBOs by Neptune. For an internal density $\rho = 2 \text{ g/cm}^3$, $M_p = M_N = 17M_\oplus$, $e_H = 0.03$, $a_p = a_d = 26.6 \text{ AU}$, $e_{\text{res}} = 0.25$, $\mathcal{M} = 2$ (so that $\Sigma_m = 0.2 \text{ g cm}^{-2}$), and $T = 3 \times 10^7 \text{ yr}$, resonant capture and retention require

$$\frac{s}{\text{km}} \lesssim \frac{s_{\text{crit}}}{\text{km}} \sim \begin{cases} 700 \mathcal{R}^{8/3} \mathcal{C}^{-2/3}, & \text{if } e < e_H/\mathcal{R}^2; \\ 70 e^{-2/3} \mathcal{R}^{4/3} \mathcal{C}^{-2/3}, & \text{if } e_H/\mathcal{R}^2 < e < \mathcal{R}e_H; \\ 7000 e^{2/3} \mathcal{C}^{-2/3}, & \text{if } e > \mathcal{R}e_H. \end{cases} \quad (53)$$

For example, if $e = 0.1$ and $\mathcal{R} = 1$, then line 3 of (53) obtains and $s_{\text{crit}} = 1500 \mathcal{C}^{-2/3} \text{ km}$. Maximum stochasticity

results when $\mathcal{R} = 1$ and $e \leq e_H$ (see also Equation [47]); either of lines 1 or 2 then yield $s_{\text{crit}} \sim 700 \mathcal{C}^{-2/3} \text{ km}$. These size estimates decrease by about 20% when corrected to reflect the fact that the width of the 3:2 resonance is somewhat smaller than the pendulum model implies (see the discussion following Equation [50]).

3.4. Analytical Formula for the Retention Fraction

As defined in §3.2, P_{keep} is the resonance retention fraction, or the probability that a typical resonant particle is retained in resonance over some duration of migration. We calculate P_{keep} by modelling the random component of the planet's migration as a diffusive continuum process. In the limit that the planet encounters a large number $\bar{N} \gg 1$ of planetesimals, the Poisson distribution (Equation [37]) is well-approximated by a Gaussian distribution with mean $\bar{N}$ and variance $\bar{N}$. Thus, over a time interval $\Delta t \gg \bar{N}^{-1}$, the random displacement of the planet, $\Delta S_{\text{rnd}} = \Delta a_p (N - \bar{N})$, has the probability density distribution

$$f(\Delta S_{\text{rnd}}, \Delta t) = \frac{1}{\sqrt{2\pi D \Delta t}} \exp(-(\Delta S_{\text{rnd}})^2 / (2D \Delta t)), \quad (54)$$

where $D = (\Delta a_p)^2 \dot{\bar{N}}$ is the diffusion coefficient and we recall that Δa_p is the change in a_p due to an encounter with a single planetesimal. The evolution of $a_{p,\text{rnd}}$ with t is continuous and the distribution f is independent over any two non-overlapping intervals Δt (the random walk has no memory). In other words, ΔS_{rnd} evolves as a Wiener process, or equivalently according to the rules of Brownian motion (e.g., Grimmer & Stirzaker 2001a). From Equation (54), it follows that over time T , the probability that $|\Delta S_{\text{rnd}}|$ does not exceed $\delta a_{p,\text{lib}}/2$ equals

$$P_{\text{keep}} = \sum_{n=1}^{\infty} \frac{4}{n\pi} \sin^3\left(\frac{n\pi}{2}\right) e^{-\lambda_n T}, \quad (55)$$

where $\lambda_n = (n\pi)^2 D / (2\delta a_{p,\text{lib}}^2)$ (see Appendix A for a derivation).

Suppose migration occurs in a disk of planetesimals having a single size s . Figure 3 displays P_{keep} as a function of s and of exponential migration timescale τ defined according to

$$a_{p,\text{avg}}(t) = a_{p,f} - (a_{p,f} - a_{p,i}) e^{-t/\tau}, \quad (56)$$

where $a_{p,i}$ and $a_{p,f}$ are the planet's initial and final average semi-major axes, respectively. In Equation (55), we take $T = 2.6\tau$, and evaluate remaining quantities for the case of maximum stochasticity: $e \leq e_H$ and $\mathcal{R} = 1$. Then $\Delta a_p = \Delta a_{p,\text{nc1}}$ (Equation [25]) and $\dot{\bar{N}} = 2\Sigma_m \Omega R_H^2/m$ (Equation [36]), with a factor of 2 inserted to account for disk material both inside and outside the planet's orbit). As in §3.2, we take $\rho = 2 \text{ g/cm}^3$, $M_p = M_N = 17M_\oplus$, $e_H = 0.03$, $a_p = a_d = 26.6 \text{ AU}$, $R_H = e_H a_p$, and $\mathcal{M} = 2$. To evaluate $\delta a_{p,\text{lib}}$, we take $e_{\text{res}} = 0.25$ for a particle in 3:2 resonance. Figure 3 describes how for a given size s , the retention fraction decreases with increasing τ ; the longer the duration of migration, the more chance a particle has of being jostled out of resonance. For $\tau = 10 \text{ Myr}$, planetesimals must have sizes $s \lesssim 500 \text{ km}$ for the retention fraction to remain greater than 1/2. These results confirm and refine our order-of-magnitude estimates

made in §3.3. Similar results were obtained for the 2:1 resonance.

The continuum limit is valid as long as the expectation value of the time required for a resonant particle to escape, $\langle t_{\text{escape}} \rangle \sim \dot{\bar{N}}^{-1} [(\delta a_{\text{p,lib}}/2)/\Delta a_{\text{p}}]^2$, greatly exceeds the time for the planet to encounter one planetesimal, $\dot{\bar{N}}^{-1}$. This criterion is satisfied for the full range of parameters adopted in Figure 3.

3.5. Numerical Results for the Retention Fraction

To explore how a stochastically migrating planet captures and retains test particles into its exterior resonances, and to test the analytic considerations of §§3.2–3.4, we perform a series of numerical integrations. We focus as before on the 3:2 (Plutino) resonance with Neptune.

Following Murray-Clay & Chiang (2005, hereafter MC05), we employ a series expansion for the time-dependent Hamiltonian,

$$H = -\frac{(GM_{\odot})^2}{2(3\Gamma + \mathcal{N})^2} - \left[\frac{GM_{\odot}}{a_{\text{p}}(t)^3} \right]^{1/2} (2\Gamma + \mathcal{N}) - \frac{GM_{\text{p}}}{a_{\text{p}}(t)} [\alpha(f_1 + f_2 e^2 + f_{31} e \cos \phi)] , \quad (57)$$

where $\alpha = a_{\text{p}}/a \approx 0.76$, the f_i 's are given in Murray & Dermott (1999), and $\mathcal{N}$ (Equation [48]) is a constant of the motion determined by initial conditions. The resonance angle,

$$\phi = 3\lambda_{\text{res}} - 2\lambda_{\text{p}} - \varpi_{\text{res}} , \quad (58)$$

is defined by the mean longitude λ_{res} and longitude of periastron ϖ_{res} of the resonant particle, and the mean longitude λ_{p} of the planet. The resonance angle librates about π for particles in resonance. The momentum conjugate to ϕ is Γ . We integrate the equations of motion,

$$\dot{\phi} = \frac{\partial H}{\partial \Gamma} , \quad \dot{\Gamma} = -\frac{\partial H}{\partial \phi} , \quad (59)$$

using the Bulirsch-Stoer algorithm (Press et al. 1992) for fixed α and f_i 's.

The Hamiltonian in Equation (57) faithfully reproduces the main features of the resonance potential; see Beaugé (1994, his Figures 12a and 12c) for a direct comparison between such a truncated Hamiltonian and the exact Hamiltonian, averaged over the synodic period (see also that paper and Murray-Clay & Chiang 2005 for a discussion of the pitfalls of keeping one too many a term in the expansion). Of course, even the exact Hamiltonian, because it is time-averaged and neglects chaotic zones, is inaccurate with regards to details such as the libration width, but these inaccuracies are slight; see the discussions following Equations (50) and (53).

To compute $a_{\text{p}}(t)$, we specify separately the average and random components of the migration velocity, $\dot{a}_{\text{p,avg}}$ and $\dot{a}_{\text{p,rand}}$. For $\dot{a}_{\text{p,avg}}$, we adopt the prescription (equivalent to Equation [56])

$$\dot{a}_{\text{p,avg}} = \frac{1}{\tau} (a_{\text{p,f}} - a_{\text{p,i}}) e^{-t/\tau} , \quad (60)$$

where $a_{\text{p,i}}$ and $a_{\text{p,f}}$ are the planet's initial and final average semi-major axes, respectively, and τ is a time constant. To compute $\dot{a}_{\text{p,rand}}$, we divide the integration into

time intervals of length $1/\Omega_{\text{p}}$. The only requirement for the time interval is that it be less than the libration period $t_{\text{lib}} \sim 400/\Omega_{\text{p}}$ (see §3.2). Over each interval, we randomly generate

$$\dot{a}_{\text{p,rand}} = \Omega_{\text{p}} \Delta a_{\text{p}} (N_{\Omega} - \dot{\bar{N}} \Omega_{\text{p}}^{-1}) . \quad (61)$$

We focus on the case of maximum stochasticity, so that $\Delta a_{\text{p}} = \Delta a_{\text{p,nc1}}$ (Equation [25]) and $\dot{\bar{N}} = 2\Sigma_m R_{\text{H}}^2 \Omega_{\text{p}}/m$ (Equation [36]) with $\mathcal{R} = 1$ and an extra factor of 2 inserted to account for disk material on both sides of the planet's orbit). We assume that the entirety of the disk mass is in planetesimals of a single mass m . Each N_{Ω} is a random deviate drawn from a Poisson distribution having mean $\dot{\bar{N}} \Omega_{\text{p}}^{-1}$.

Figure 4 displays the sample evolution of a test particle driven into 3:2 resonance by a stochastically migrating planet. For this integration, $M_{\text{p}} = M_{\text{N}}$, $a_{\text{p,i}} = 23.1$ AU, $a_{\text{p,f}} = 30.1$ AU, $\tau = 10^7$ yr, $\mathcal{R} = 1$, $\mathcal{M} = 2$ (so that $\Sigma_m = 0.2 \text{ g cm}^{-2}$), and $s = 150$ km ($m = 3 \times 10^{22}$ g). This choice for s is sufficiently small that the particle is successfully captured and retained by the planet.

Contrast Figure 4 with Figure 5, in which all model parameters are the same except for a larger $s = 700$ km. In this case the planet eventually loses the test particle because the migration is too noisy.

Figure 6 displays the fraction of particles caught and kept in resonance as a function of τ and s . For each data point in Figure 6, we follow the evolution of 200 particles initialized with eccentricities of approximately 0.01 and semi-major axes that lie outside the initial position of resonance by about 1 AU.⁸ Figure 6 is the numerical counterpart of Figure 3; the agreement between the two is excellent and validates our analytic considerations. If $\tau = 10^7$ yr (consistent with findings by MC05), then the capture fraction rises above 0.5 for $s \lesssim 500$ km. Since these results pertain to the case $\{\mathcal{R} = 1, e \leq e_{\text{H}}\}$ which yields the largest amount of noise for given Σ_m and s , we conclude that $s \sim 500$ km is the lowest, and thus the most conservative, estimate we can make for the maximum planetesimal size compatible with resonant capture of KBOs by a migrating Neptune, assuming that the entire disk is composed of planetesimals of a single size (this assumption is relaxed in the next section). In other words, if Neptune's migration were driven by planetesimals all having $s \ll 500$ km, stochasticity would not have impeded the trapping of Resonant KBOs. Of course, our numerical estimate of 500 km is uncertain insofar as we have not kept track of order-unity constants in our derivations. We suspect a more careful analysis will revise our size estimate downwards by a factor of a few (see the discussion of $\mathcal{C}$ in §2.3.3).

3.6. Planetesimal Size Distributions

Actual disks comprise planetesimals with a range of sizes. From Equation (46), the stochasticity in the planet's migration is dominated by those planetesimals having maximal $\Sigma_m m$. What was the distribution of sizes during the era of Neptune's migration? A possible answer is provided by the coagulation simulations

⁸ The particles do not all have the same initial eccentricities and semi-major axes. This is because they occupy the same Hamiltonian level curve; see section 3.5 of MC05.




FIG. 3.— Fraction of particles retained in external 3:2 resonance by a stochastically migrating planet as a function of migration timescale, calculated according to Equation (55). The entire disk mass is assumed to be in planetesimals of a single size s , and a range of choices for s are shown. The diffusivity D is evaluated at its maximum value, appropriate for the case $e \leq e_H$ and $\mathcal{R} = 1$. We set $\Delta a_p = \Delta a_{p,nc1}$ (Equation [25]), $\tilde{N} = 2\Sigma_m \Omega R_H^2 / m$, (Equation [36]), $\rho = 2 \text{ g/cm}^3$, $M_p = M_N = 17M_\oplus$, $e_H = 0.03$, $a_p = a_d = 26.6 \text{ AU}$, $R_H = e_H a_p$, $\mathcal{M} = 2$, $e_{\text{res}} = 0.25$, and $T = 2.6\tau$. Planetesimals having sizes smaller than $\sim 200 \text{ km}$ produce so little noise in the planet's migration that no object is lost from the 3:2 resonance. Compare this Figure with its numerical counterpart, Figure 6. In calculating P_{keep} , we assume $\mathcal{C} = 1$; probably $\mathcal{C}$ is of order several, in which case the sizes indicated in the Figure should be revised downward by a factor of a few ($\mathcal{C}^{2/3}$; see Equation [53]).

of Kenyon & Luu (1999, hereafter KL99). The left-hand panel of their Figure 8 portrays the evolution of the size distribution, starting with a disk of seed bodies having sizes up to 100 m and a total surface density of $\Sigma = 0.2 \text{ g cm}^{-2}$. After $t = 11 \text{ Myr}$, the size bin for which $\Sigma_m m$ is maximal is centered at $s \sim 4 \text{ km}$; for this bin at that time, $\Sigma_m = 10^{-3} \text{ g cm}^{-2}$ (evaluated within a logarithmic size interval 0.3 dex wide). After $t = 37 \text{ Myr}$, the planetesimals generating the most stochasticity have $s \sim 750 \text{ km}$ and $\Sigma_m = 2 \times 10^{-3} \text{ g cm}^{-2}$. Note that at $t = 37 \text{ Myr}$, the stochasticity is dominated by the largest planetesimals formed, but they do not contain the bulk of the total disk mass; the lion's share of the mass is instead sequestered into km-sized objects.

In Figure 7, we plot the resonance retention fraction P_{keep} (Equation [55]) for the KL99 size distribution at $t = 11$ and 37 Myr , using the values of $s(m)$ and Σ_m cited above. The remaining parameters that enter into P_{keep} are chosen to be the same as those employed for Figure 3; i.e., we adopt the case of maximum stochasticity. Evidently, $P_{\text{keep}} = 1$ for the KL99 size distributions; stochasticity is negligible.

For comparison, we also plot in Figure 7 the retention fraction for pure power-law size distributions: $d\eta/ds \propto s^{-q}$, where $d\eta$ is the differential number of planetesimals having sizes between s and $s + ds$. Since $\Sigma_m m \propto s^{7-q}$, stochasticity is dominated by the upper end of the size

distribution for $q < 7$. We fix the maximal radius to be that of Pluto ($s_{\text{upper}} = 1200 \text{ km}$), set the total surface density $\Sigma = 0.2 \text{ g cm}^{-2}$, and calculate P_{keep} for three choices of $q = 3.5, 4$, and 4.5 . For $q \geq 4$, the lower limit of the size distribution significantly influences the normalization of $d\eta/ds$; for $q = 4$ and 4.5 , we experiment with two choices for the minimum planetesimal radius, $s_{\text{lower}} = 1 \text{ km}$ and 1 m . We equate Σ_m with the integrated surface density between $s_{\text{upper}}/2$ and s_{upper} . According to Figure 7, steep size distributions $q \geq 4$ are characterized by order-unity retention efficiencies. In contrast, shallow size distributions $q < 4$ for which the bulk of the mass is concentrated towards s_{upper} can introduce significant stochasticity.

4. CONCLUDING REMARKS

We summarize our findings in §4.1 and discuss quantitatively some remaining issues in §4.2.

4.1. Summary

Newly formed planets likely occupy remnant planetesimal disks. Planets migrate as they exchange energy and angular momentum with planetesimals. Driven by discrete scattering events, migration is stochastic.

In our solar system, Neptune may have migrated outward by several AU and thereby captured the many Kuiper belt objects (KBOs) found today in mean-motion




FIG. 4.— Evolution of a particle caught into 3:2 resonance with a stochastically migrating planet. Stochasticity is driven by a disk of surface density $\Sigma_m = 0.2 \text{ g cm}^{-2}$, all in planetesimals having sizes $s = 150 \text{ km}$ and sub-Hill random velocities. The random walk in the planet’s semi-major axis causes the libration amplitude of the resonant particle to undergo a corresponding random walk. The noise in this example is too mild to prevent the planet from both capturing and retaining the particle in resonance.




FIG. 5.— Evolution of a particle caught into, but eventually lost from, 3:2 resonance with a stochastically migrating planet. Stochasticity is driven by a disk of surface density $\Sigma_m = 0.2 \text{ g cm}^{-2}$, all in planetesimals having sizes $s = 700 \text{ km}$ and sub-Hill random velocities. The particle is expelled from resonance having had its eccentricity raised to 0.2 during its time in resonant lock.

resonance with the planet. While resonance capture is efficient when migration is smooth, a longstanding issue has been whether Neptune’s actual migration was too noisy to permit capture. Our work addresses—and dispels—this concern by supplying a first-principles theory for how a planet’s semi-major axis fluctuates in response to intrinsic granularity in the gravitational potential. We apply our theory to identify the environmental conditions under which resonance capture remains viable.

Stochasticity results from random variations in the numbers and orbital properties of planetesimals encountering the planet. The degree of stochasticity (as measured, say, by $\sigma_{a_p, T}$, the typical distance that the planet’s semi-major axis random walks away from its average value) depends on how planetesimal semi-major axes a and random velocities u are distributed. We have parameterized a by its difference from the planet’s semi-major axis: $x \equiv a - a_p \equiv \mathcal{R}R_H$, where R_H is the Hill sphere radius and $\mathcal{R} \gtrsim 1$. In the case of high dispersion when

$u > \mathcal{R}v_H$ (where $v_H \equiv \Omega_p R_H$ is the Hill velocity and Ω_p is the planet’s orbital angular velocity), planetesimal orbits cross that of the planet. Stochasticity increases with decreasing u in the high-dispersion case because the cross-section for strong scatterings increases steeply with decreasing velocity dispersion (as $1/u^4$). In the intermediate-dispersion case when $v_H/\mathcal{R}^2 < u < \mathcal{R}v_H$, planetesimal and planet orbits do not cross, and stochasticity decreases with decreasing u . In the low-dispersion case when $u < v_H/\mathcal{R}^2$, the amount of stochasticity is insensitive to u .

The values of u and $\mathcal{R}$ which actually characterize disks are unknown. The random velocity u , for example, is expected to be set by a balance between excitation by gravitational scatterings and damping by inelastic collisions between planetesimals and/or gas drag. Damping depends, in turn, on the size distribution of planetesimals. These considerations are often absent from current N-body simulations of planetary migration in planetesimal disks. Despite such uncertainty, we can still identify the circumstances under which stochasticity is maximal.




FIG. 6.— Fraction of particles caught into, and retained within, external 3:2 resonance by a stochastically migrating planet. For every τ and s , we numerically integrate the trajectories of 200 test particles with initial eccentricities of ~ 0.01 and semi-major axes that lie 1 AU outside of nominal resonance. These particles respond to the time-averaged potential of a Neptune-mass planet which migrates outward from 23.1 AU to 30.1 AU within a disk of fixed surface density $\Sigma_m = 0.2 \text{ g cm}^{-2}$ in planetesimals of a single size s . The planetesimals have sub-Hill random velocities and semi-major axes that lie within $\mathcal{R} = 1$ Hill radius of the planet's; these choices maximize the amount of stochasticity in the planet's migration. Compare this Figure with its analytic counterpart, Figure 3; the agreement is excellent. The solid curve labelled “Smooth” corresponds to the case when all noise is eliminated from the planet's migration. Planetesimals having sizes smaller than ~ 200 km yield an essentially smooth migration. For $\tau \lesssim 10^5$ yr, capture is not possible even if migration were smooth, since the migration is too fast to be adiabatic. These results are calculated for $C = 1$; probably C is of order several and so the sizes indicated in the Figure should be revised downward by a factor of a few ($C^{2/3}$; see Equation [53]).

Maximum stochasticity obtains when $\mathcal{R} \sim 1$ and $u \lesssim v_H$, that is, when planetesimals have semi-major axes within a Hill radius of the planet's and when their velocity dispersion is no greater than the Hill velocity.

A stochastically migrating planet cannot retain objects in a given resonance if the planet's semi-major axis random walks away from its average value by a distance greater than the maximum libration width of the resonance. This simple criterion is validated by numerical experiments and enables analytic calculation of the resonance retention efficiency as a function of disk parameters. A disk of given surface density generates more noise when composed of fewer, larger planetesimals. In the context of Neptune's migration, we estimate that if the bulk of the minimum-mass disk resided in bodies having sizes smaller than $\mathcal{O}(100)$ km and if the fraction of the disk mass in larger bodies was not too large ($\lesssim$ a few percent for planetesimals having sizes of 1000 km, for example), then the retention efficiency of Neptune's first-order resonances would have been of order unity ($\gtrsim 0.1$). Such order-unity efficiencies seem required by observations, which *prima facie* place $122/474 \approx 26\%$ of well-observed KBOs (excluding Centaurs) inside mean-motion resonances (Chiang et al. 2006). Drawing conclusions based on a comparison between this observed percentage and our theoretical retention percentage P_{keep}

is a task fraught with caveats—a more fair comparison would require, e.g., disentangling the observational bias against discovering Resonant vs. non-Resonant objects; account of the attrition of the Resonant population due to weak chaos over the four-billion-year age of the solar system; and knowledge of the initial eccentricity and semi-major axis distributions of objects prior to resonance sweeping, as these distributions impact capture probabilities in different ways for different resonances (Chiang et al. 2003; Hahn & Malhotra 2005; Chiang et al. 2006). But each of these caveats alters the relevant percentages only by factors of a few, and when combined, their effects tend to cancel. Therefore we feel comfortable in our assessment that P_{keep} must have been of order unity to explain the current Resonant population. In that case, $\mathcal{O}(100)$ km is a conservative estimate for the maximum allowed size of planetesimals comprising the bulk of the disk mass, derived for the case of maximum stochasticity.

How does an upper limit of $\mathcal{O}(100)$ km compare with the actual size distribution of the planetesimal disk? While today's Kuiper belt places most of its mass in objects having sizes of ~ 100 km, this total mass is tiny—only $\sim 0.1 M_{\oplus}$ (Bernstein et al. 2004; see Chiang et al. 2006 for a synopsis). The current belt is therefore 2–3 orders of magnitude too low in mass to have driven Nep-




FIG. 7.— Fraction of particles retained in external 3:2 resonance by a stochastically migrating planet for various planetesimal size distributions. The retention efficiency is calculated analytically using Equation (55), with parameters the same as those for Figure 3 except for $\Sigma_m \times m$; that parameter is evaluated at its maximum value within a logarithmic size bin spanning a factor of 2 for a given size distribution. The size distributions considered include two from Kenyon & Luu (1999; their Figure 8), evaluated at times $t = 11$ Myr and 37 Myr; and five different power-law distributions, each characterized by a total integrated surface density $\Sigma = 0.2 \text{ g cm}^{-2}$, an upper size limit $s_{\text{upper}} = 1200 \text{ km}$, a differential size index q (such that $d\eta/ds \propto s^{-q}$), and a lower size limit s_{lower} as indicated (the curve for $q = 3.5$ is insensitive to s_{lower} since the bulk of the mass is concentrated towards s_{upper}). The three curves for the size distributions of KL99 and for $\{q = 4.5, s_{\text{lower}} = 1 \text{ m}\}$ overlap at $P_{\text{keep}} = 1$.

tune’s migration. The current size distribution is such that bodies having radii $\gtrsim 40 \text{ km}$ are collisionless over the age of the solar system and might therefore represent a direct remnant, unadulterated by erosive collisions, of the planetesimal disk during the era of migration (Pan & Sari 2005). If so, the bulk of the primordial disk mass must have resided in bodies having sizes $\lesssim 40 \text{ km}$. Theoretical calculations of the coagulation history of the Kuiper belt are so far consistent with this expectation. Kenyon & Luu (1999) find, for their primordial trans-Neptunian disk of $10M_{\oplus}$, that 99% of the mass failed to coagulate into bodies larger than $\mathcal{O}(1) \text{ km}$, because the formation of several Pluto-sized objects (comprising $\sim 0.1\%$ of the total mass) excited velocity dispersions so much that planetesimal collisions became destructive rather than agglomerative. The average-mass planetesimals in their simulation have sizes $\mathcal{O}(1) \text{ km}$, much smaller than even our most conservative estimate of the maximum allowed size of $\mathcal{O}(100) \text{ km}$.

For a given size distribution of planetesimals, most stochasticity is produced by the size bin having maximal ηm^2 , which need not be the size bin containing the majority of the mass. Here, η and m are the number of planetesimals and the mass of an individual planetesimal in a logarithmic size bin. For power-law size distributions $d\eta/ds \propto s^{-q}$ such that $q < 7$, stochasticity is dominated by the largest planetesimals. For disks having

as much mass as the minimum-mass disk of solids and whose largest members are Pluto-sized, size distributions with $q \geq 4$ enjoy order-unity efficiencies for resonance retention. The size distributions of Kenyon & Luu (1999) resemble $q = 4$ power laws, but with a large overabundance of planetesimals having sizes of $\mathcal{O}(1) \text{ km}$. This sequestration of mass dramatically reduces the stochasticity generated by the largest bodies, which have sizes of $\mathcal{O}(1000) \text{ km}$.

We conclude that Neptune’s Brownian motion did not impede in any substantive way the planet’s capture and retention of Resonant KBOs.

4.2. Extensions

4.2.1. Single Kick to Planet

Our focus thus far has been on the regime in which many stochastic kicks to the planet are required for resonant particles to escape. Of course, a single kick from a planetesimal having sufficiently large mass m_1 could flush particles from resonance. To estimate m_1 , we equate the change in the planet’s semi-major axis from a single encounter, Δa_p , to the maximum half-width of the resonance, $\delta a_{p,\text{lib}}/2$ (see Equation [50] and related discussion). In the likely event that the perturber’s eccentricity e is of order unity, then $\max(\Delta a_p) \sim (m_1/M_p)a_p e$ (Equation [31]) and therefore $m_1 \gtrsim 0.6 (0.5/e) M_{\oplus}$ for Plutinos to escape resonance. Our estimate for m_1 agrees

with that of Malhotra (1993). Why such enormous perturbations have not been observed today is unclear and casts doubt on their existence (Morbidelli, Jacob, & Petit 2002). If such an Earth-mass planetesimal were present over the duration T of Neptune’s migration, then the likelihood of a resonance-destabilizing encounter would be $P_1 \sim \dot{N}T \sim 10^{-2}(0.5/e)^4$, where the encounter rate $\dot{N}$ is given by Equation (42) with $\Sigma_m \sim m_1/(2\pi a_p^2)$, and we have set $T \sim 2.7 \times 10^7$ yr.

4.2.2. Kicks to Resonant Planetesimals

Finally, we have ignored in this work how disk planetesimals directly perturb the semi-major axis of a resonant particle. This neglect does not significantly alter our conclusions. Take the resonant planetesimal to resemble a typical Resonant KBO observed today, having size $s_{\text{res}} \sim 100$ km. Then its Hill velocity is $e_{\text{H,res}}\Omega a \sim 10^{-4}\Omega a$. The relative velocity between the resonant planetesimal and an ambient, perturbing planetesimal greatly exceeds this Hill velocity, if only because migration in resonant lock quickly raises the eccentricity of the resonant planetes-

imal above $e_{\text{H,res}}$. Equation (43), appropriate for the super-Hill regime, implies that $\langle \dot{a}_{\text{p,rd}}^2 \rangle^{1/2} \propto M_p^0$ —the RMS random velocity does not depend on the mass of the object being perturbed! Therefore when both the resonant planetesimal and the planet are scattering planetesimals in the super-Hill regime, their random walks are comparable in vigor. The conservative limit of $\mathcal{O}(100)$ km on the planetesimal size which allows resonance retention is derived, by contrast, for the sub-Hill, maximum stochasticity regime, and is therefore little affected by these considerations.⁹

This work was made possible by grants from the National Science Foundation, NASA, and the Alfred P. Sloan Foundation. We thank B. Collins, J. Hahn, A. Morbidelli, I. Shapiro, J. Wisdom, and an anonymous Protostars and Planets V referee for encouraging and informative exchanges. Section 3.6 was written in response to insightful comments by Mike Brown and Re’em Sari. An anonymous referee helped to improve the presentation of this paper.

⁹ The probability that a planetesimal will be ejected from resonance by a planetesimal of comparable mass in a single encounter

is negligibly small.

APPENDIX

ABSORPTION PROBABILITY FOR BROWNIAN MOTION WITH A DOUBLE BOUNDARY

Here we derive Equation (55), the probability that a particle experiencing Brownian motion between two absorbing boundaries has not been absorbed by time t (e.g., Grimmett and Stirzaker 2001b). Consider Brownian motion along a path $x(t)$ with $x(0) = 0$ and absorbing boundaries at $x = \pm b$, $b \geq 0$. The probability density distribution $f(x, t)$ satisfies the diffusion equation

$$\frac{\partial f}{\partial t} = \frac{1}{2}D \frac{\partial^2 f}{\partial x^2}, \quad (\text{A1})$$

where D is the diffusion coefficient. The absorbing boundaries generate the boundary conditions

$$f(\pm b, t) = 0 \quad (\text{A2})$$

for all time,¹⁰ and the initial condition is

$$f(x, 0) = \delta(x). \quad (\text{A3})$$

To solve for $f(x, t)$, we expand f in a Fourier series, keeping only terms that satisfy (A2):

$$f(x, t) = \sum_{n=1}^{\infty} k_n(t) \sin\left(\frac{n\pi(x+b)}{2b}\right). \quad (\text{A4})$$

Plugging (A4) into (A1), we find

$$k_n(t) = c_n e^{-\lambda_n t}, \quad (\text{A5})$$

where the c_n ’s are constants and

$$\lambda_n \equiv \frac{n^2\pi^2}{8b^2}D. \quad (\text{A6})$$

The c_n ’s must satisfy (A3). From Fourier analysis at time $t = 0$, we find

$$c_n = \frac{1}{2b} \int_0^{4b} [\delta(y-b) - \delta(y-3b)] \sin\left(\frac{n\pi y}{2b}\right) dy \quad (\text{A7})$$

$$= \frac{1}{b} \sin\left(\frac{n\pi}{2}\right). \quad (\text{A8})$$

¹⁰ Equation (A2) holds as long as D is non-zero. Particles near the boundary are carried across by fluctuations too quickly to maintain a non-zero density f at $x = \pm b$ (see Grimmett and Stirzaker 2001a for a proof).

The probability that the walker has not yet crossed either of the absorbing boundaries at time t is

$$P_{\text{keep}}(t) = \int_{-b}^b f(x, t) dx \tag{A9}$$

$$= \int_{-b}^b \sum_{n=1}^{\infty} c_n e^{-\lambda_n t} \sin\left(\frac{n\pi(x+b)}{2b}\right) dx \tag{A10}$$

$$= \sum_{n=1}^{\infty} \frac{4}{n\pi} \sin^3\left(\frac{n\pi}{2}\right) e^{-(n\pi)^2 Dt/(8b^2)} . \tag{A11}$$

REFERENCES

- Chiang, E.I. 2003, *ApJ*, 584, 465
 Chiang, E.I., et al. 2003, *AJ*, 126, 430
 Chiang, E.I., et al. 2006, in *Protostars and Planets V*, eds. B. Reipurth, K. Keil, & D. Jewitt (Tucson: Univ. Arizona Press), in press
 Collins, B.F. & Sari, R. 2006, *AJ*, in press
 Cruikshank, D., et al. 2006, in *Protostars and Planets V*, eds. B. Reipurth, K. Keil, & D. Jewitt (Tucson: Univ. Arizona Press), in press
 Dermott, S.F., Malhotra, R., & Murray, C.D. 1988, *Icarus*, 76, 295
 Dones, L., Weissman, P.R., Levison, H.F., & Duncan, M.J. 2004, *ASPC Proceedings*, 323, 371
 Fernández, J.A., & Ip, W.H. 1984, *Icarus*, 58, 109
 Franklin, F.A., et al. 2004, *AJ*, 128, 1391
 Goldreich, P. 1965, *MNRAS*, 130, 159
 Goldreich, P., Lithwick, Y., & Sari, R. 2004, *ARA&A*, 42, 549
 Gomes, R.S., Morbidelli, A., & Levison, H.F. 2004, *Icarus*, 170, 492
 Grimmett, G., & Stirzaker, D. 2001a, *Probability and Random Processes* (3rd ed.; Oxford: Oxford Univ. Press)
 Grimmett, G., & Stirzaker, D. 2001b, *One Thousand Exercises in Probability* (Oxford: Oxford Univ. Press)
 Hahn, J.M., & Malhotra, R. 1999, *AJ*, 117, 3041
 Hahn, J.M., & Malhotra, R. 2000, *DPS Meeting #32*, #19.06
 Hahn, J.M., & Malhotra, R. 2005, *AJ*, 130, 2392
 Hénon, M., & Petit, J.M. 1986, *Celestial Mechanics*, 38, 67
 Ida, S., & Makino, J. 1992, *Icarus*, 96, 107
 Kenyon, S.J., & Luu, J.X. 1999, *AJ*, 118, 1101
 Laughlin, G., Steinacker, A., & Adams, F.C. 2004, *ApJ*, 608, 489
 Lee, M.H., et al. 2006, *ApJ*, 641, L1178
 Levison, H.F., & Morbidelli, A. 2003, *Nature*, 426, 419
 Levison, H.F., Morbidelli, A., Gomes, R.S., & Backman, D. 2006, in *Protostars and Planets V*, eds. B. Reipurth, K. Keil, & D. Jewitt (Tucson: Univ. Arizona Press), in press
 Malhotra, R. 1993, *Nature*, 365, 819
 Malhotra, R. 1995, *AJ*, 110, 420
 Meyer, M., et al. 2006, in *Protostars and Planets V*, eds. B. Reipurth, K. Keil, & D. Jewitt (Tucson: Univ. Arizona Press), in press
 Morbidelli, A., Brown, M.E., & Levison, H.F. 2003, *Earth Moon & Planets*, 92, 1
 Morbidelli, A., Jacob, C., & Petit, J.M. 2002, *Icarus*, 157, 241
 Murray, C.D., & Dermott, S.F. 1999, *Solar System Dynamics* (Cambridge: Cambridge Univ. Press)
 Murray, N., et al. 1998, *Science*, 279, 69
 Murray-Clay, R.A., & Chiang, E.I. 2005, *ApJ*, 619, 623 (MC05)
 Nelson, R.P. 2005, *A&A*, 443, 1067
 Peale, S.J. 1986, in *Satellites*, eds. J.A. Burns & M.S. Matthews (Tucson: Univ. Arizona Press), 159
 Petit, J.M., & Hénon, M. 1986, *Icarus*, 66, 536
 Quillen, A.C. 2006, *MNRAS*, 365, 1367
 Rafikov, R.R. 2003, *AJ*, 126, 2529
 Trujillo, C.A., & Brown, M.E. 2001, *ApJ*, 554, 95
 Tsiganis, K., Gomes, R., Morbidelli, A., & Levison, H.F. 2005, *Nature*, 435, 459
 Vogt, S., et al. 2005, *ApJ*, 632, 638
 Wisdom, J. 1980, *AJ*, 85, 1122
 Wyatt, M.C. 2003, *ApJ*, 598, 1321
 Zhou, L., et al. 2002, *MNRAS*, 336, 520

TABLE A1
FREQUENTLY USED SYMBOLS

Symbol	Definition	Remark
t	Time	...
a	Planetesimal Semi-Major Axis	...
e	Planetesimal Eccentricity	...
Ω	Planetesimal Orbital Angular Velocity	...
u	Planetesimal Random (Epicyclic) Velocity	$\sim e\Omega a$
m	Planetesimal Mass	...
s	Planetesimal Radius	...
ρ	Planetesimal Internal Density	2 g cm^{-3}
a_d	Mean Radius of Planetesimal Disk Annulus	...
Σ	Total Disk Surface Density (Mass Per Unit Face-On Area)	0.2 g cm^{-2} for minimum-mass trans-Neptunian disk
Σ_m	Disk Surface Density in Planetesimals of Mass m	...
$\mathcal{M}$	$2\pi\Sigma_m a_d^2/M_p$, Parameterizes Surface Density	2 for $\Sigma_m = 0.2 \text{ g cm}^{-2}$, $a_d = 26.6 \text{ AU}$, and $M_p = M_N$
a_p	Planet Semi-Major Axis	...
Ω_p	Planet Orbital Angular Velocity	...
M_p	Planet Mass	...
M_*	Mass of Host Star	...
R_H	Hill Radius of Planet $\equiv a_p(M_p/(3M_*))^{1/3}$	...
e_H	Hill Eccentricity $\equiv R_H/a_p$	...
v_H	Hill Velocity $\equiv \Omega_p R_H$	...
$\dot{a}_{p,\text{rnd}}$	Planet Random Migration Velocity	time-averages to zero
$\dot{a}_{p,\text{avg}}$	Planet Average Migration Velocity	assumed known function
x	$a - a_p$	$ x \lesssim a_p$
$\mathcal{R}$	Minimum Value of $ x /R_H$	$\gtrsim 1$
b	Impact Parameter of Planet-Planetesimal Encounter	> 0
Δt_e	Duration of Planet-Planetesimal Encounter	$\sim 1/\Omega_p$ at longest
ΔQ	Change in Quantity Q from a Single Encounter	evaluated well before and well after encounter, e.g., Δa_p
Δt	Arbitrary Time Interval	...
N	Number of Planetesimals Encountered by Planet in Δt	Poisson deviate
$\bar{N}$	Mean of N	...
$\bar{\dot{N}}$	Mean Rate of Planetesimal Encounters by Planet	...
$\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2}$	Root-Mean-Squared (RMS) Random Migration Velocity Over Δt	$\propto 1/\sqrt{\Delta t}$
$\mathcal{C}$	Numerical Coefficient for $\langle \dot{a}_{p,\text{rnd}}^2 \rangle^{1/2}$	Equation (46), estimated to be of order several
D	Diffusivity of Planet's Semi-Major Axis $= (\Delta a_p)^2 \bar{\dot{N}}$	...
$a_{p,i}$	Initial Semi-Major Axis of Planet, Pre-Migration	23.1 AU
$a_{p,f}$	Final Semi-Major Axis of Planet, Post-Migration	30.1 AU
τ	Exponential Timescale for Migration	Equation (56)
T	Total Duration of Migration	...
$\sigma_{a_p,T}$	$\langle (a_p - a_{p,\text{avg}})^2 \rangle^{1/2}$ After Time T	$\propto T^{1/2}$, Equation (51)
P_{keep}	Probability a Resonant Particle is Retained in Resonance After Time T	Equation (55)
S_{rnd}	$\int_0^t \dot{a}_{p,\text{rnd}} dt$	...
ΔS_{rnd}	$S_{\text{rnd}}(t + \Delta t) - S_{\text{rnd}}(t)$	...
m_{crit}	Maximum Planetesimal Mass Satisfying $P_{\text{keep}} \sim 1$, For Disks of a Single Planetesimal Mass	Equation (52)
s_{crit}	Maximum Planetesimal Radius Satisfying $P_{\text{keep}} \sim 1$, For Disks of a Single Planetesimal Mass	Equation (53)
f	Probability Density	Equation (54)
M_N	Mass of Neptune	...
e_{res}	Eccentricity of Resonant Planetesimal	...
ϕ	Resonance Angle (Libration Phase) of Resonant Planetesimal	Equation (58)
$\delta a_{p,\text{lib}}$	Maximum Width of Resonance, Referred to Planet's Orbit	Equation (50) and related discussion
$d\eta/ds$	Differential Size Spectrum of Noise-Generating Planetesimals	...
q	Index for Power-Law Size Distributions	$d\eta/ds \propto s^{-q}$